

• **Name:** Tetsuro SUGIURA

• **Current Position:**

Director, Kansai Medical University Medical Center, Japan
Professor, Kochi Medical School, Japan

• **Country:** Japan

• **Educational Background:**

- MD, Kansai Medical University, Japan (1976)
- PhD, Kansai Medical University, Japan (1982)

• **Professional Experience:**

- Professor, Kochi Medical School(1999~present)
- Director, Kochi Medical School Hospital(2010~present)
- Director, Kansai Medical University Medical Center(2018~present)
- Chairman, 12nd Cherry Blossom Symposium in Awaji, Hyogo, Japan(2022)

• **Professional Organizations:**

- Japanese Society of Internal Medicine(Fellow)
- Japanese Society of Cardiology(Fellow)
- Japanese College of Cardiology(Fellow)
- American College of Chest Physicians(Fellow)
- Japanese Society of Laboratory Medicine(Fellow)

• **Main Scientific Publications:**

Jiao YF, Okumiya T, Saibara T, Kudo Y, Sugiura T.

Erythrocyte creatine as a marker of excessive erythrocyte destruction due to hypersplenism in patients with liver cirrhosis. *Clin Biochem.* 34:395-8,2001

Kudo Y, Yamasaki F, Doi Y, Sugiura T.

Clinical correlates of PR-segment depression in asymptomatic patients with pericardial effusion. *J Am Coll Cardiol.* 39:2000-4, 2002

Kudo Y, Yamasaki F, Doi T, Doi Y, Sugiura T.

Clinical significance of low voltage in asymptomatic patients with pericardial effusion free of heart disease. *Chest.* 124:2064-7, 2003

Okumiya T, Ishikawa-Nishi M, Doi T, Kamioka M, Takeuchi H, Doi Y, Sugiura T. Evaluation of intravascular hemolysis with erythrocyte creatine in patients with cardiac valve prostheses. *Chest*. 125:2115-20, 2004

Sugiura T, Nakamura S, Kudo Y, Okumiya T, Yamasaki F, Iwasaka T. Clinical factors associated with persistent pericardial effusion after successful primary coronary angioplasty. *Chest*. 128:798-803, 2005

Con SA, Takeuchi H, Valerin AL, Con-Wong R, Con-Chin GR, Con-Chin VG, Nishioka M, Mena F, Brenes F, Yasuda N, Araki K, Sugiura T. Diversity of *Helicobacter pylori* cagA and vacA Genes in Costa Rica: its Relationship with Atrophic Gastritis and Gastric Cancer. *Helicobacter*. 12:547-52, 2007

Morishita S, Nishimori I, Minakuchi T, Onishi S, Takeuchi H, Sugiura T, Vulio D, Scozzafava A. Supuran CT. Cloning, polymorphism, and inhibition of beta-carbonic anhydrase of *Helicobacter pylori*. *J Gastroenterol*. 43:849-57, 2008

Kubo T, Kitaoka H, Okawa M, Yamanaka S, Hirota T, Baba Y, Hayato K, Yamasaki N, Matsumura Y, Yasuda N, Sugiura T, Doi YL. Combined measurements of cardiac troponin I and brain natriuretic peptide are useful for predicting adverse outcomes in hypertrophic cardiomyopathy. *Circ J*. 75:919 -26, 2011

Kubo T, Kitaoka H, Yamanaka S, Hirota T, Baba Y, Hayashi K, Iiyama T, Kumagai N, Tanioka K, Yamasaki N, Matsumura Y, Furuno T, Sugiura T, Doi YL. Significance of High-Sensitivity Cardiac Troponin T in Hypertrophic Cardiomyopathy. *J Am Coll Cardiol*. 62(8):1252-9, 2013

Sugiura T, Yamanaka S, Takeuchi H, Morimoto N, Kamioka M, Matsumura Y. Autoimmunity and pulmonary hypertension in patients with Graves' disease. *Heart Vessels*. 30(6):642-6, 2015

Sugiura T, Okuyama T, Kamioka M, Kubo T, Hirakawa Y, Hisahara T, Matsumura Y. Intravascular hemolysis in patients with mitral regurgitation: Evaluation by erythrocyte creatine. *J cardiol*. 71:414-8, 2018

Matsumura K, Okumiya T, Sugiura T, Takahashi N, Yamamoto Y, Kikuchi S, Fujii K, Otagaki M, Shiojima I. Shortened red blood cell age in patients with end-stage renal disease who were receiving haemodialysis: a cross-sectional study. *BMC Nephrol*. 21:418-24, 2020